The Plant Journal (2000) **24**(6), 905–910

Author index

Abbadi, A., Brummel, M. and Spener, F. Knockout of the regulatory site of 3-ketoacyl-ACP synthase III enhances short- and medium-chain acyl-ACP synthesis 1
Abbo, S. *see* Gorbunova, V.
Abranches, R., Santos, A.P., Wegel, E., Williams, S., Castilho, A., Christou, P., Shaw, P. and Stoger, E. Widely separated multiple transgene integration sites in wheat chromosomes are brought together at interphase 713
Allan, A.C. *see* Wood, N.T.
Allan, V. *see* Barroso, C.
Alvarez-Buylla, E.R., Liljegren, S.J., Pelaz, S., Gold, S.E., Burgeff, C., Ditta, G.S., Vergara-Silva, F. and Yanofsky, M.F. MADS-box gene evolution beyond flowers: expression in pollen, endosperm, guard cells, roots and trichomes 457
Amano, Y. *see* Ohmiya, Y.
Angenent, G.C. *see* Ge, Y.-X.
Aoki, N. *see* Bauer, C.S.
Aoyama, T. *see* Sakai, H.
Apel, K. *see* Borner, R.
Aquilani, R. *see* Rosati, C.
Arazi, T. *see* Sunkar, R.
Atkinson, H. *see* Urwin, P.
Ausubel, F.M. *see* Dewdney, J.
Avivi-Ragolski, N. *see* Gorbunova, V.

Baba, T. *see* Hase, Y.
Backhaus, R.A. *see* Bouvier, F.
Bagnaresi, P., Mazars-Marty, D., Pupillo, P., Marty, F. and Briat, J.-F. Tonoplast subcellular localization of maize cytochrome b_5 reductases 645
Balagué, C. *see* Godard, F.
Bao, F. *see* Hu, Y.
Barroso, C., Chan, J., Allan, V., Doonan, J., Hussey, P. and Lloyd,C. Two kinesin-related proteins associated with the cold-stable cytoskeleton of carrot cells: characterization of a novel kinesin, DcKRP120-2 859
Bauer, C.S., Hoth, S., Haga, K., Philippar, K., Aoki, N. and Hedrich,R. Differential expression and regulation of K^+ channels in the maize coleoptile: molecular and biophysical analysis of cells isolated from cortex and vasculature 139
Baur, B. *see* Häusler, R.E.
Bechtold, N. *see* Sarrobert, C.
Bechtold, N. *see* Sherson, S.M.
Beck, C.F. *see* Kropat, J.
Bernier, G. *see* Bonhomme, F.
Besenbeck, R. *see* Meyer, S.
Bhalerao, R.P. *see* Yamaguchi, M.
Blatt, M.R. *see* Leyman, B.
Blumwald, E. *see* Snedden, W.A.
Bogucki, A. *see* Hanin, M.
Bohnert, H.J. *see* Chauhan, S.
Bollini, R. *see* Sparvoli, F.
Bonhomme, F., Kurz, B., Melzer, S., Bernier, G. and Jacqmard, A. Cytokinin and gibberellin activate *SaMADS A*, a gene apparently involved in the regulation of the floral transition in *Sinapis alba* 103
Borner, R., Kampmann, G., Chandler, J., Gleißner, R., Wisman, E., Apel, K. and Melzer, S. A MADS domain gene involved in the transition to flowering in *Arabidopsis* 591
Bouché, N. *see* Sunkar, R.
Bouchez, D. *see* Sunkar, R.
Bougourd, S., Marrison, J. and Haseloff, J. An aniline blue staining procedure for confocal microscopy and 3D imaging of normal and perturbed cellular phenotypes in mature *Arabidopsis* embryos 543
Bouvier, F. *see* Rosati, C.
Bouvier, F., Suire, C., d'Harlingue, A., Backhaus, R.A. and Camara, B. Molecular cloning of geranyl diphosphate synthase and compartmentation of monoterpene synthesis in plant cells 241
Bowyer, J.R. *see* Vinti, G.
Brady, S.R. *see* Hu, S.
Bramley, P.M. *see* Fraser, P.D.
Bray, C.M. *see* West, C.E.
Brennicke, A. *see* Okada, S.
Brennicke, A. *see* Schock, I.
Briat, J.-F. *see* Bagnaresi, P.
Brummel, M. *see* Abbadi, A.
Burgeff, C. *see* Alvarez-Buylla, E.R.
Busconi, L. *see* Martín, M.L.
Busscher, M. *see* Ge, Y.-X.

Caldelari, D. *see* Rodriguez-Concepción, M.
Camara, B. *see* Bouvier, F.
Camara, B. *see* Rosati, C.
Campbell, S. *see* Vinti, G.
Castilho, A. *see* Abranches, R.
Ceriotti, A. *see* Sparvoli, F.
Chan, J. *see* Barroso, C.
Chandler, J. *see* Borner, R.
Chang, K.S., Lee, S.H., Hwang, S.B. and Park, K.Y. Characterization and translational regulation of the arginine decarboxylase gene in carnation (*Dianthus caryophyllus* L.) 45
Chatfield, S.P., Stirnberg, P., Forde, B.G. and Leyser, O. The hormonal regulation of axillary bud growth in *Arabidopsis* 159
Chauhan, S., Forsthoefel, N., Ran, Y., Quigley, F., Nelson, D.E. and Bohnert, H.J. Na^+/*myo*-inositol symporters and Na^+/H^+-antiport in *Mesembryanthemum crystallinum* 511
Chen, Z. *see* Du, L.
Chételat, A. *see* Vollenweider, S.
Chevalier, T. *see* Marin-Olivier, M.
Chory, J. *see* Vinti, G.
Christou, P. *see* Abranches, R.
Chua, N.-H. *see* Zuo, J.
Clark, G.B. *see* Hu, S.
Clarke, A., Desikan, R., Hurst, R.D., Hancock, J.T. and Neill, S.J. NO way back: nitric oxide and programmed cell death in *Arabidopsis thaliana* suspension cultures 667
Contard-David, P. *see* Sarrobert, C.
Creemers-Molenaar, T. *see* Ge, Y.-X.
Criqui, M.C., Parmentier, Y., Derevier, A., Shen, W.-H., Dong, A. and Genschik, P. Cell cycle-dependent proteolysis and ectopic overexpression of cyclin B1 in tobacco BY2 cells 763
Crouch, N.P. *see* Irmler, S.
Cui, J. *see* Dewdney, J.
Cushman, J.C. *see* Patharkar, O.R.

d'Harlingue, A. *see* Bouvier, F.
Dahlhaus, E. *see* Ge, Y.-X.
Daminati, M.G. *see* Sparvoli, F.
Davies, H.V. *see* Taylor, M.A.

© 2000 Blackwell Science Ltd

de la Peña, A. *see* Martin, A.C.
De Roover, J. *see* Van den Ende, W.
de Wit, P.J.G.M. *see* Takken, F.L.W.
Deák, M. *see* Oberschall, A.
del Pozo, J.C. *see* Martin, A.C.
Derevier, A. *see* Criqui, M.C.
Desikan, R. *see* Clarke, A.
Dessi, P., Rudhe, C. and Glaser, E. Studies on the topology of the protein import channel in relation to the plant mitochondrial processing peptidase integrated into the cytochrome bc_1 complex 637
Devoto, A. *see* Dewdney, J.
Dewdney, J., Reuber, T.L., Wildermuth, M.C., Devoto, A., Cui, J., Stutius, L.M., Drummond, E.P. and Ausubel, F.M. Three unique mutants of *Arabidopsis* identify *eds* loci required for limiting growth of a biotrophic fungal pathogen 205
Dharmapuri, S. *see* Rosati, C.
Ditta, G.S. *see* Alvarez-Buylla, E.R.
Dolev, D. *see* Sunkar, R.
Dong, A. *see* Criqui, M.C.
Doonan, J. *see* Barroso, C.
Drummond, E.P. *see* Dewdney, J.
Du, L. and Chen, Z. Identification of genes encoding receptor-like protein kinases as possible targets of pathogen- and salicylic acid-induced WRKY DNA-binding proteins in *Arabidopsis* 837
Dudits, D. *see* Oberschall, A.
Dudler, R. *see* Schweizer, P.
Dumas, C. *see* Marin-Olivier, M.
Duncan, G. *see* Taylor, M.A.

Ehtesham, N.Z. *see* Pham, X.H.
Eicks, M. *see* Häusler, R.E.
Ellerström, M. *see* Ezcurra, I.
Ezcurra, I., Wycliffe, P., Nehlin, L., Ellerström, M. and Rask, L. Transactivation of the *Brassica napus* napin promoter by ABI3 requires interaction of the conserved B2 and B3 domains of ABI3 with different *cis*-elements: B2 mediates activation through an ABRE, whereas B3 interacts with an RY/G-box 57

Fabian, T. *see* Yamaguchi, M.
Faoro, F. *see* Sparvoli, F.
Farmer, E.E. *see* Vollenweider, S.
Fehér, A. *see* Oberschall, A.
Felle, H.H., Hanstein, S., Steinmeyer, R. and Hedrich, R. Dynamics of ionic activities in the apoplast of the sub-stomatal cavity of intact *Vicia faba* leaves during stomatal closure evoked by ABA and darkness 297
Finnemann, J. and Schjoerring, J.K. Post-translational regulation of cytosolic glutamine synthetase by reversible phosphorylation and 14-3-3 protein interaction 171
Fischer, K. *see* Häusler, R.E.
Fischer, K.L. *see* Häusler, R.E.
Flügge, U.-I. *see* Häusler, R.E.
Fobis-Loisy, I. *see* Marin-Olivier, M.
Forbes, S. *see* Sherson, S.M.
Forde, B.G. *see* Chatfield, S.P.
Forsthoefel, N. *see* Chauhan, S.
Fowke, L.C. *see* Wang, H.
Fowler, J.E. *see* Ivanchenko, M.
Franken, J. *see* Ge, Y.-X.
Fraser, P.D., Pinto, M.E.S., Holloway, D.E. and Bramley, P.M. Application of high-performance liquid chromatography with photodiode array detection to the metabolic profiling of plant isoprenoids 551
Fromm, H. *see* Sunkar, R.
Fujisawa, M. *see* Okada, S.
Fukuzawa, H. *see* Okada, S.
Furbank, R. *see* Hurry, V.
Fuse, T. *see* Horiguchi, G.

Gabriëls, S.H.E.J. *see* Takken, F.L.W.
Gaude, T. *see* Marin-Olivier, M.
Ge, Y.-X., Angenent, G.C., Wittich, P.E., Peters, J., Franken, J., Busscher, M., Zhang, L.-M., Dahlhaus, E., Kater, M.M., Wullems, G.J. and Creemers-Molenaar, T. *NEC1*, a novel gene, highly expressed in nectary tissue of *Petunia hybrida* 725
Geelen, D. *see* Leyman, B.
Genschik, P. *see* Criqui, M.C.
Germain, V. *see* Sherson, S.M.
Giedt, C.D. and Weil, C.F. The maize *LAG1-O* mutant suggests that reproductive cell lineages show unique gene expression patterns early in vegetative development 815
Gilmartin, P.M. *see* Urwin, P.
Gilmer, S. *see* Wang, H.
Gineste, S. *see* Sarrobert, C.
Giuliano, G. *see* Rosati, C.
Glaser, E. *see* Dessi, P.
Gleißner, R. *see* Borner, R.
Godard, F., Lummerzheim, M., Saindrenan, P., Balagué, C. and Roby, D. *hxc2*, an *Arabidopsis* mutant with an altered hypersensitive response to *Xanthomonas campestris* pv. *campestris* 749
Gold, S.E. *see* Alvarez-Buylla, E.R.
Golombek, S. *see* Weber, H.
Gorbunova, V., Avivi-Ragolski, N., Shalev, G., Kovalchuk, I., Abbo, S., Hohn, B. and Levy, A.A. A new hyperrecombinogenic mutant of *Nicotiana tabacum* 601
Grant, J.J., Yun, B.-W. and Loake, G.J. Oxidative burst and cognate redox signalling reported by luciferase imaging: identification of a signal network that functions independently of ethylene, SA and Me-JA but is dependent on MAPKK activity 569
Gregan, J. *see* Schock, I.
Gruissem, W. *see* Rodríguez-Concepción, M.
Gubatz, S. *see* Weber, H.

Haga, K. *see* Bauer, C.S.
Haley, A. *see* Wood, N.T.
Hall, M. *see* Turner, S.R.
Hamada, S., Onouchi, H., Tanaka, H., Kudo, M., Liu, Y.-G., Shibata, D., Machida, C. and Machida, Y. Mutations in the *WUSCHEL gene of Arabidopsis thaliana* result in the development of shoot without juvenile leaves 91
Hancock, J.T. *see* Clarke, A.
Hanin, M., Mengiste, T., Bogucki, A. and Paszkowski, J. Elevated levels of intrachromosomal homologous recombination in *Arabidopsis* overexpressing the *MIM* gene 183
Hanstein, S. *see* Felle, H.H.
Hase, Y., Tanaka, A., Baba, T. and Watanabe, H. *FRL1* is required for petal and sepal development in *Arabidopsis* 21
Haseloff, J. *see* Bougourd, S.
Hause, B., Stenzel, I., Miersch, O., Maucher, H., Kramell, R., Ziegler, J. and Wasternack, C. Tissue-specific oxylipin signature of tomato flowers: allene oxide cyclase is highly expressed in distinct flower organs and vascular bundles 113
Häusler, R.E., Baur, B., Scharte, J., Teichmann, T., Eicks, M., Fischer, K.L., Flügge, U.-I., Schubert, S., Weber, A. and

Fischer, K. Plastidic metabolite transporters and their physiological functions in the inducible crassulacean acid metabolism plant *Mesembryanthemum crystallinum* 285
Hayashi, T. *see* Ohmiya, Y.
Hedrich, R. *see* Bauer, C.S.
Hedrich, R. *see* Felle, H.H.
Heim, U. *see* Weber, H.
Hemmann, G. *see* Sherson, S.M.
Hills, A. *see* Vinti, G.
Hohn, B. *see* Gorbunova, V.
Holloway, D.E. *see* Fraser, P.D.
Horiguchi, G., Fuse, T., Kawakami, N., Kodama, H. and Iba, K. Temperature-dependent translational regulation of the ER ω-3 fatty acid desaturase gene in wheat root tips 805
Horváth, G.V. *see* Oberschall, A.
Hoth, S. *see* Bauer, C.S.
Hotze, M. *see* Irmler, S.
Howell, S.H. *see* Mockaitis, K.
Hu, S., Brady, S.R., Kovar, D.R., Staiger, C.J., Clark, G.B., Roux, S.J. and Muday, G.K. Identification of plant actin-binding proteins by F-actin affinity chromatography 127
Hu, Y., Bao, F. and Li, J. Promotive effect of brassinosteroids on cell division involves a distinct *CycD3*-induction pathway in *Arabidopsis* 693
Hümmer, C. *see* Meyer, S.
Hurry, V., Strand, Å., Furbank, R. and Stitt, M. The role of inorganic phosphate in the development of freezing tolerance and the acclimatization of photosynthesis to low temperature is revealed by the *pho* mutants of *Arabidopsis thaliana* 383
Hurst, R.D. *see* Clarke, A.
Hussey, P. *see* Barroso, C.
Hwang, S.B. *see* Chang, K.S.

Iba, K. *see* Horiguchi, G.
Ichimura, K., Mizoguchi, T., Yoshida, R., Yuasa, T. and Shinozaki, K. Various abiotic stresses rapidly activate *Arabidopsis* MAP kinases ATMPK4 and ATMPK6 655
Iglesias, J. *see* Martin, A.C.
Iida, A., Kazuoka, T., Torikai, S., Kikuchi, H. and Oeda, K. A zinc finger protein RHL41 mediates the light acclimatization response in *Arabidopsis* 191
Irmler, S., Schröder, G., St-Pierre, B., Crouch, N.P., Hotze, M., Schmidt, J., Strack, D., Matern, U. and Schröder, J. Indole alkaloid biosynthesis in *Catharanthus roseus*: new enzyme activities and identification of cytochrome P450 CYP72A1 as secologanin synthase 797
Ivanchenko, M., Vejlupkova, Z., Quatrano, R.S. and Fowler, J.E. Maize ROP7 GTPase contains a unique, CaaX box-independent plasma membrane targeting signal 79

Jacqmard, A. *see* Bonhomme, F.
Jiang, Q. *see* West, C.E.
Joosten, M.H.A.J. *see* Takken, F.L.W.

Kampmann, G. *see* Borner, R.
Kanda, T. *see* Ohmiya, Y.
Kaplan, B. *see* Sunkar, R.
Kater, M.M. *see* Ge, Y.-X.
Kawakami, N. *see* Horiguchi, G.
Kazuoka, T. *see* Iida, A.
Kendrick, R.E. *see* Weller, J.L.
Kikuchi, H. *see* Iida, A.
Kirschner, M., Winkelhaus, S., Thierfelder, J.M. and Nover, L. Transient expression and heat-stress-induced co-aggregation of endogenous and heterologous small heat-stress proteins in tobacco protoplasts 397
Knoop, V. *see* Schock, I.
Kodama, H. *see* Horiguchi, G.
Kono, K. *see* Okada, S.
Koornneef, M. *see* Weller, J.L.
Koprek, T., McElroy, D., Louwerse, J., Williams-Carrier, R. and Lemaux, P.G. An efficient method for dispersing *Ds* elements in the barley genome as a tool for determining gene function 253
Kovács, I. *see* Oberschall, A.
Kovalchuk, I. *see* Gorbunova, V.
Kovar, D.R. *see* Hu, S.
Kovar, D.R., Staiger, C.J., Weaver, E.A. and McCurdy, D.W. AtFim1 is an actin filament crosslinking protein from *Arabidopsis thaliana* 625
Kramell, R. *see* Hause, B.
Kropat, J., Oster, U., Rüdiger, W. and Beck, C.F. Chloroplast signalling in the light induction of nuclear *HSP70* genes requires the accumulation of chlorophyll precursors and their accessibility to cytoplasm/nucleus 523
Kudo, M. *see* Hamada, S.
Kurz, B. *see* Bonhomme, F.

Lee, S.H. *see* Chang, K.S.
Lemaux, P.G. *see* Koprek, T.
Levy, A.A. *see* Gorbunova, V.
Leyman, B., Geelen, D. and Blatt, M.R. Localization and control of expression of Nt-Syr1, a tobacco snare protein 369
Leyser, O. *see* Chatfield, S.P.
Leyva, A. *see* Martin, A.C.
Li, J. *see* Hu, Y.
Li, J. *see* Ouyang, J.
Liljegren, S.J. *see* Alvarez-Buylla, E.R.
Liu, Y.-G. *see* Hamada, S.
Lloyd, C. *see* Barroso, C.
Loake, G.J. *see* Grant, J.J.
López-Juez, E. *see* Vinti, G.
Louwerse, J. *see* Koprek, T.
Lu, R. *see* Takken, F.L.W.
Lucas, W.J. *see* Xoconostle-Cázares, B.
Luderer, R. *see* Takken, F.L.W.
Ludwig, A., Stolz, J. and Sauer, N. Plant sucrose-H^+ symporters mediate the transport of vitamin H 503
Lummerzheim, M. *see* Godard, F.

Maathuis, F.J.M. *see* Sunkar, R.
Machida, C. *see* Hamada, S.
Machida, Y. *see* Hamada, S.
Marin-Olivier, M., Chevalier, T., Fobis-Loisy, I., Dumas, C. and Gaude, T. Aquaporin *PIP* genes are not expressed in the stigma papillae in *Brassica oleracea* 231
Marrison, J. *see* Bougourd, S.
Martin, A.C., del Pozo, J.C., Iglesias, J., Rubio, V., Solano, R., de la Peña, A., Leyva, A. and Paz-Ares, J. Influence of cytokinins on the expression of phosphate starvation responsive genes in *Arabidopsis* 559
Martin, H. *see* Urwin, P.
Martín, M.L. and Busconi, L. Membrane localization of a rice calcium-dependent protein kinase (CDPK) is mediated by myristoylation and palmitoylation 429

Marty, F. *see* Bagnaresi, P.
Marusic, C. *see* Rosati, C.
Matern, U. *see* Irmler, S.
Matta, B. *see* Pham, X.H.
Maucher, H. *see* Hause, B.
Mazars-Marty, D. *see* Bagnaresi, P.
McCurdy, D.W. *see* Kovar, D.R.
McElroy, D. *see* Koprek, T.
McRae, D. *see* Taylor, M.A.
Melzer, M. *see* Meyer, S.
Melzer, S. *see* Bonhomme, F.
Melzer, S. *see* Borner, R.
Mengiste, T. *see* Hanin, M.
Meyer, S., Melzer, M., Truernit, E., Hümmer, C., Besenbeck, R., Stadler, R. and Sauer, N. *AtSUC3*, a gene encoding a new *Arabidopsis* sucrose transporter, is expressed in cells adjacent to the vascular tissue and in a carpel cell layer 869
Michiels, A. *see* Van den Ende, W.
Miersch, O. *see* Hause, B.
Millam, S. *see* Taylor, M.A.
Mizoguchi, T. *see* Ichimura, K.
Mochizuki, N. *see* Vinti, G.
Mockaitis, K. and Howell, S.H. Auxin induces mitogenic activated protein kinase (MAPK) activation in roots of Arabidopsis seedlings 785
Muday, G.K. *see* Hu, S.

Nakayama, S. *see* Okada, S.
Nehlin, L. *see* Ezcurra, I.
Neill, S.J. *see* Clarke, A.
Nelson, D.E. *see* Chauhan, S.
Nishiyama, R. *see* Okada, S.
Niu, Q.-W. *see* Zuo, J.
Nover, L. *see* Kirschner, M.
Nussaume, L. *see* Sarrobert, C.

Oberschall, A., Deák, M., Török, K., Sass, L., Vass, I., Kovács, I., Fehér, A., Dudits, D. and Horváth, G.V. A novel aldose/aldehyde reductase protects transgenic plants against lipid peroxidation under chemical and drought stresses 437
Oeda, K. *see* Iida, A.
Ohmiya, Y., Samejima, M., Shiroishi, M., Amano, Y., Kanda, T., Sakai, F. and Hayashi, T. Evidence that endo-1,4-β-glucanases act on cellulose in suspension-cultured poplar cells 147
Ohyama, K. *see* Okada, S.
Oka, A. *see* Sakai, H.
Okada, S., Fujisawa, M., Sone, T., Nakayama, S., Nishiyama, R., Takenaka, M., Yamaoka, S., Sakaida, M., Kono, K., Takahama, M., Yamato, K.T., Fukuzawa, H., Brennicke, A. and Ohyama, K. Construction of male and female PAC genomic libraries suitable for identification of Y-chromosome-specific clones from the liverwort, *Marchantia polymorpha* 421
Onouchi, H. *see* Hamada, S.
Oster, U. *see* Kropat, J.
Ouyang, J., Shao, X. and Li, J. Indole-3-glycerol phosphate, a branchpoint of indole-3-acetic acid biosynthesis from tryptophan biosynthetic pathway in *Arabidopsis thaliana* 327

Pallara, P. *see* Rosati, C.
Park, K.Y. *see* Chang, K.S.
Parmentier, Y. *see* Criqui, M.C.
Paszkowski, J. *see* Hanin, M.
Patharkar, O.R. and Cushman, J.C. A stress-induced calcium-dependent protein kinase from *Mesembryanthemum crystallinum* phosphorylates a two-component pseudo-response regulator 679
Paz-Ares, J. *see* Martin, A.C.
Pelaz, S. *see* Alvarez-Buylla, E.R.
Peters, J. *see* Ge, Y.-X.
Pham, X.H., Reddy, M.K., Ehtesham, N.Z., Matta, B. and Tuteja, N. A DNA helicase from *Pisum sativum* is homologous to translation initiation factor and stimulates topoisomerase I activity 219
Philippar, K. *see* Bauer, C.S.
Pinto, M.E.S. *see* Fraser, P.D.
Pokorny, J. *see* Schweizer, P.
Pupillo, P. *see* Bagnaresi, P.

Quatrano, R.S. *see* Ivanchenko, M.
Quigley, F. *see* Chauhan, S.

Ran, Y. *see* Chauhan, S.
Rask, L. *see* Ezcurra, I.
Reddy, M.K. *see* Pham, X.H.
Reuber, T.L. *see* Dewdney, J.
Robaglia, C. *see* Sarrobert, C.
Roberts, I. *see* Taylor, M.A.
Roby, D. *see* Godard, F.
Rodríguez-Concepción, M., Toledo-Ortiz, G., Yalovsky, S., Caldelari, D. and Gruissem, W. Carboxyl-methylation of prenylated calmodulin CaM53 is required for efficient plasma membrane targeting of the protein 775
Rolletschek, H. *see* Weber, H.
Rosati, C., Aquilani, R., Dharmapuri, S., Pallara, P., Marusic, C., Tavazza, R., Bouvier, F., Camara, B. and Giuliano, G. Metabolic engineering of beta-carotene and lycopene content in tomato fruit 413
Ross, H.A. *see* Taylor, M.A.
Roux, S.J. *see* Hu, S.
Rubio, V. *see* Martin, A.C.
Rudhe, C. *see* Dessi, P.
Rüdiger, W. *see* Kropat, J.
Ruiz-Medrano, R. *see* Xoconostle-Cázares, B.

Saindrenan, P. *see* Godard, F.
Sakai, F. *see* Ohmiya, Y.
Sakai, H., Aoyama, T. and Oka, A. *Arabidopsis* ARR1 and ARR2 response regulators operate as transcriptional activators 703
Sakaida, M. *see* Okada, S.
Samejima, M. *see* Ohmiya, Y.
Sandberg, G. *see* Yamaguchi, M.
Sanders, D. *see* Sunkar, R.
Santos, A.P. *see* Abranches, R.
Sarrobert, C., Thibaud, M.-C., Contard-David, P., Gineste, S., Bechtold, N., Robaglia, C. and Nussaume, L. Identification of an *Arabidopsis thaliana* mutant accumulating threonine resulting from mutation in a new dihydrodipicolinate synthase gene 357
Sass, L. *see* Oberschall, A.
Sauer, N. *see* Ludwig, A.
Sauer, N. *see* Meyer, S.
Sauer, N. *see* Sherson, S.M.
Sauter, M. *see* Yamaguchi, M.
Scharte, J. *see* Häusler, R.E.
Schjoerring, J.K. *see* Finnemann, J.

Schmidt, J. *see* Irmler, S.
Schock, I., Gregan, J., Steinhauser, S., Schweyen, R., Brennicke, A. and Knoop, V. A member of a novel *Arabidopsis thaliana* gene family of candidate Mg^{2+} ion transporters complements a yeast mitochondrial group II intron-splicing mutant 489
Schrader, J. *see* Yamaguchi, M.
Schreuder, M.E.L. *see* Weller, J.L.
Schröder, G. *see* Irmler, S.
Schröder, J. *see* Irmler, S.
Schubert, S. *see* Häusler, R.E.
Schulze-Lefert, P. *see* Schweizer, P.
Schweizer, P., Pokorny, J., Schulze-Lefert, P. and Dudler, R. Double-stranded RNA interferes with gene function at the single-cell level in cereals 895
Schweyen, R. *see* Schock, I.
Shalev, G. *see* Gorbunova, V.
Shao, X. *see* Ouyang, J.
Shaw, P. *see* Abranches, R.
Shen, W.-H. *see* Criqui, M.C.
Sherson, S.M., Hemmann, G., Wallace, G., Forbes, S., Germain, V., Stadler, R., Bechtold, N., Sauer, N. and Smith, S.M. Monosaccharide/proton symporter AtSTP1 plays a major role in uptake and response of *Arabidopsis* seeds and seedlings to sugars 849
Shibata, D. *see* Hamada, S.
Shinozaki, K. *see* Ichimura, K.
Shiroishi, M. *see* Ohmiya, Y.
Smith, H. *see* Weller, J.L.
Smith, S.M. *see* Sherson, S.M.
Snedden, W.A. and Blumwald, E. Alternative splicing of a novel diacylglycerol kinase in tomato leads to a calmodulin-binding isoform 317
Solano, R. *see* Martin, A.C.
Sone, T. *see* Okada, S.
Sparvoli, F., Faoro, F., Daminati, M.G., Ceriotti, A. and Bollini, R. Misfolding and aggregation of vacuolar glycoproteins in plant cells 825
Spener, F. *see* Abbadi, A.
Stadler, R. *see* Meyer, S.
Stadler, R. *see* Sherson, S.M.
Staiger, C.J. *see* Hu, S.
Staiger, C.J. *see* Kovar, D.R.
Steinhauser, S. *see* Schock, I.
Steinmeyer, R. *see* Felle, H.H.
Stenzel, I. *see* Hause, B.
Stewart, D. *see* Taylor, M.A.
Stirnberg, P. *see* Chatfield, S.P.
Stitt, M. *see* Hurry, V.
Stoger, E. *see* Abranches, R.
Stolz, J. *see* Ludwig, A.
Stolz, S. *see* Vollenweider, S.
St-Pierre, B. *see* Irmler, S.
Strack, D. *see* Irmler, S.
Strand, Å. *see* Hurry, V.
Stutius, L.M. *see* Dewdney, J.
Suire, C. *see* Bouvier, F.
Sunkar, R., Kaplan, B., Bouché, N., Arazi, T., Dolev, D., Talke, I.N., Maathuis, F.J.M., Sanders, D., Bouchez, D. and Fromm, H. Expression of a truncated tobacco *NtCBP4* channel in transgenic plants and disruption of the homologous *Arabidopsis* CNGC1 gene confer Pb^{2+} tolerance 533

Takahama, M. *see* Okada, S.
Takenaka, M. *see* Okada, S.
Takken, F.L.W., Luderer, R., Gabriëls, S.H.E.J., Westerink, N., Lu, R., de Wit, P.J.G.M. and Joosten, M.H.A.J. A functional cloning strategy, based on a binary PVX-expression vector, to isolate HR-inducing cDNAs of plant pathogens 275
Talke, I.N. *see* Sunkar, R.
Tanaka, A. *see* Hase, Y.
Tanaka, H. *see* Hamada, S.
Tavazza, R. *see* Rosati, C.
Taylor, M.A., Ross, H.A., McRae, D., Stewart, D., Roberts, I., Duncan, G., Wright, F., Millam, S. and Davies, H.V. A potato α-glucosidase gene encodes a glycoprotein-processing α-glucosidase II-like activity. Demonstration of enzyme activity and effects of down-regulation in transgenic plants 305
Teichmann, T. *see* Häusler, R.E.
Thibaud, M.-C. *see* Sarrobert, C.
Thierfelder, J.M. *see* Kirschner, M.
Toledo-Ortiz, G. *see* Rodríguez-Concepción, M.
Torikai, S. *see* Iida, A.
Török, K. *see* Oberschall, A.
Trewavas, A.J. *see* Wood, N.T.
Truernit, E. *see* Meyer, S.
Turner, S.R. and Hall, M. The *gapped xylem* mutant identifies a common regulatory step in secondary cell wall deposition 477
Tuteja, N. *see* Pham, X.H.

Uchimiya, H. *see* Yamaguchi, M.
Umeda, M. *see* Yamaguchi, M.
Urwin, P., Yi, L., Martin, H., Atkinson, H. and Gilmartin, P.M. Functional characterization of the EMCV IRES in plants 583

Van den Ende, W., Michiels, A., De Roover, J., Verhaert, P. and Van Laere, A. Cloning and functional analysis of chicory root fructan 1-exohydrolase I (1-FEH I): a vacuolar enzyme derived from a cell-wall invertase ancestor? Mass fingerprint of the 1-FEH I enzyme 447
Van Laere, A. *see* Van den Ende, W.
Vass, I. *see* Oberschall, A.
Vejlupkova, Z. *see* Ivanchenko, M.
Vergara-Silva, F. *see* Alvarez-Buylla, E.R.
Verhaert, P. *see* Van den Ende, W.
Vinti, G., Hills, A., Campbell, S., Bowyer, J.R., Mochizuki, N., Chory, J. and López-Juez, E. Interactions between *hy1* and *gun* mutants of *Arabidopsis*, and their implications for plastid/nuclear signalling 883
Viry-Moussaïd, M. *see* Wood, N.T.
Vollenweider, S., Weber, H., Stolz, S., Chételat, A. and Farmer, E.E. Fatty acid ketodienes and fatty acid ketotrienes: Michael addition acceptors that accumulate in wounded and diseased *Arabidopsis* leaves 467

Wallace, G. *see* Sherson, S.M.
Wang, H., Zhou, Y., Gilmer, S., Whitwill, S. and Fowke, L.C. Expression of the plant cyclin-dependent kinase inhibitor ICK1 affects cell division, plant growth and morphology 613
Wasternack, C. *see* Hause, B.
Watanabe, H. *see* Hase, Y.
Waterworth, W.M. *see* West, C.E.
Weaver, E.A. *see* Kovar, D.R.
Weber, A. *see* Häusler, R.E.
Weber, H. *see* Vollenweider, S.
Weber, H., Rolletschek, H., Heim, U., Golombek, S., Gubatz, S. and Wobus, U. Antisense-inhibition of ADP-glucose pyrophosphorylase in developing seeds of *Vicia narbonensis* moderately decreases starch but increases protein content and affects seed maturation 33

Wegel, E. *see* Abranches, R.
Weil, C.F. *see* Giedt, C.D.
Weller, J.L., Schreuder, M.E.L., Smith, H., Koornneef, M. and Kendrick, R.E. Physiological interactions of phytochromes A, B1 and B2 in the control of development in tomato 345
West, C.E., Waterworth, W.M., Jiang, Q. and Bray, C.M. *Arabidopsis* DNA ligase IV is induced by γ-irradiation and interacts with an *Arabidopsis* homologue of the double strand break repair protein XRCC4 67
Westerink, N. *see* Takken, F.L.W.
Whitwill, S. *see* Wang, H.
Wildermuth, M.C. *see* Dewdney, J.
Williams, S. *see* Abranches, R.
Williams-Carrier, R. *see* Koprek, T.
Winkelhaus, S. *see* Kirschner, M.
Wisman, E. *see* Borner, R.
Wittich, P.E. *see* Ge, Y.-X.
Wobus, U. *see* Weber, H.
Wood, N.T., Allan, A.C., Haley, A., Viry-Moussaïd, M. and Trewavas, A.J. The characterization of differential calcium signalling in tobacco guard cells 335
Wright, F. *see* Taylor, M.A.
Wullems, G.J. *see* Ge, Y.-X.
Wycliffe, P. *see* Ezcurra, I.

Xoconostle-Cázares, B., Ruiz-Medrano, R. and Lucas, W.J. Proteolytic processing of CmPP36, a protein from the cytochrome b_5 reductase family, is required for entry into the phloem translocation pathway 735

Yalovsky, S. *see* Rodríguez-Concepción, M.
Yamaguchi, M., Fabian, T., Sauter, M., Bhalerao, R.P., Schrader, J., Sandberg, G., Umeda, M. and Uchimiya, H. Activation of CDK-activating kinase is dependent on interaction with H-type cyclins in plants 11
Yamaoka, S. *see* Okada, S.
Yamato, K.T. *see* Okada, S.
Yanofsky, M.F. *see* Alvarez-Buylla, E.R.
Yi, L. *see* Urwin, P.
Yoshida, R. *see* Ichimura, K.
Yuasa, T. *see* Ichimura, K.
Yun, B.-W. *see* Grant, J.J.

Zhang, L.-M. *see* Ge, Y.-X.
Zhou, Y. *see* Wang, H.
Ziegler, J. *see* Hause, B.
Zuo, J., Niu, Q.-W. and Chua, N.-H. An estrogen receptor-based transactivator XVE mediates highly inducible gene expression in transgenic plants 265

The Plant Journal (2000) **24**(6), 911–914

Subject index

α-glucosidase 305
α-linolenic acid 805
ABA 57, 297, 335, 369
ABI3 57
abiotic stress 679
abiotic stress (salt, wounding) 369
ABRE 57
abscisic acid 159
Ac 815
Ac transposase 253
actin cytoskeleton 127
actin filament crosslinking 625
ADP-glucose pyrophosphorylase 33
AGL20 591
Agrobacterium tumefaciens 275
aldo-keto reductase 437
allene oxide cyclase 113
aniline blue 543
annexin 127
anther stomium 725
anthocyanin 895
antisense 305, 413
antisense RNA 327
apical dominance 159
apoplast 297
aquaporin 231
Arabidopsis 21, 205, 357, 383, 457, 467, 477, 489, 503, 511, 559, 591, 625, 837
Arabidopsis thaliana 91, 103, 183, 327, 543, 613, 693, 749, 849
arginine decarboxylase 45
aspartate 357
AtFim1 625
auxin 147, 159
auxin-activated gene expression 785
auxin response mutants 785
auxin transport 127

b_5 reductase 645
barley 253, 895
bc_1 complex 637
BimC 859
biotin transport 503
Blumeria graminis 895
Brassica 231
Brassica napus L. 171, 725
brassinosteroid 693
BRCT domain 67
bud growth 159

calcium 533, 679
calcium-dependent protein kinase 429, 679
calcium signalling 335
calmodulin 317, 775
CAM 285
CaM53 775
carnation 45
carrot cells 859
caspase 667
Catharanthus roseus 797
CDK-activating kinase 11
CDK inhibitor 613
cell cycle 11, 763
cell death 477
cell differentiation 33
cell division 613, 693
cell size 613
cell-to-cell trafficking 735
cellular architecture 543
cellulose 147
cGMP 667
chlorophyll precursor pools 523
chromatin organization 713
Cichorium intybus 447
Cladosporium fulvum 275
cold acclimatization 383
cold shock 335
complementation cloning 503
confocal microscopy 543, 713
Cucurbita maxima 735
Cuphea 1
CycD3 693
cyclic nucleotides 533
cyclin 763
cyclin-dependent protein kinase 11
cyclin H 11
cytochrome b_5 reductase 735
cytochrome P450 797
cytokinin 103, 159, 559
cytoskeleton 625
cytosol 241

3D visualization 543
Dbox 763
DEAD-box protein 219
de-etiolation 345
defence-related 895
defense response 205
detoxification 437
DHDPS 357
diacylglycerol 317
dioecious plant 421
disease susceptibility 205
diurnal cycle 285
DNA binding 703
DNA repair 67, 183
DNA replication 219
drought tolerance 437
Ds-transposition 253

elicitor 275
embryo 543
encephalomyocarditis virus (ECMV) 583
endo,1-4-β-glucanase 147
endomembrane 775
endoplasmic reticulum 825
endo-reduplication 21
endosperm 457
epidermal strips 335
Erysiphe orontii 205
estradiol-inducible 265
estrogen receptor 265
ethylene 159

© 2000 Blackwell Science Ltd

exocytosis 369
exporter 869
expression 503, 869

F-actin affinity chromatography 127
fatty acid synthase 1
1-FEH 447
fimbrin/plastin 625
FISH 713
floral induction 591
floral signal 103
flower 21
fluorescence *in situ* hybridization (FISH) 421
frl1 21
fructan breakdown 447
function 583
functional genomics 253

gene expression 265
gene family 489
gene tagging 253
gene targeting 601
geranyl diphosphate synthase 241
germination 849
GFP 79
gibberellin 103
glucose transporter 285
glucosidase II 305
glutamine synthetase 171
glycoprotein processing 305
glycosylation 825
group II intron splicing cofactors 489
guard cells 335, 457
gun mutants 883

heat-stress granules 397
heavy metals 533
high light acclimatization 191
high-performance liquid chromatography 551
homologous recombination 183
hormone signaling 785
HSP70 light induction 523
hy1 mutant 883
hyper-osmolarity 655
hyperrecombinogenic mutants 601
hypersensitive response 275, 569, 749

IAA biosynthesis 327
in situ RT-PCR 735
indole 327
indole acetic acid 785
indole alkaloids 797
indole-3-glycerol phosphate 327
insertional mutagen 253
internal ribosome entry site (IRES) 583
intracellular communication 883
intracellular transport 825
inulin 447
ion channel 533
ion-selective micro-electrodes 297
isoprenoids 551

jam mutant 91
jasmonates 113
juvenile leaves 91

K^+ channel 139
3-ketoacyl-ACP synthase III 1
kinesin-like protein 859
knockout mutant 533

leaf identity 91
legume seed development 33
LexA 265
lignin 477
lipid 467, 805
lipid peroxidation 437
localization signal 79
loganin 797
low temperature 383, 655, 805
lycopene beta-cyclase 413
lysine 357

MADS box 591, 457
magnesium homeostasis 489
maize coleoptile 139
maize 79, 645
MALDI-TOF 447
malonaldehyde 467
MAP kinase 655, 667
Marchantia polymorpha 421
mechanical 335
medium-chain fatty acids 1
MEK inhibitor 785
membrane localization 429
meristem 815
Mesembryanthemum 511
Mesembryanthemum crystallinum 285, 679
metabolic profiling 551
metabolic regulation 33
metal reductase 645
Michael addition 467
micro-injection 625
microtubule-associated proteins (MAPs) 859
MIM 183
mitochondrial processing peptidase 637
Mlo 895
monosaccharide transporter 849
monoterpene compartmentation 241
mutant 205, 345, 357, 477, 749, 849
Myb 703
myristoylation 429

naphthylphthalamic acid 127
napin 57
NEC1 gene 725
nectar secretion 725
nectary 725
Nicotiana 369
Nicotiana tabacum 601
nitric oxide 667
nitrogen remobilization 171
non-homologous end joining 67

ω-3 fatty acid desaturase 805
organ development 21
overexpression 413
ovules 113
oxidative burst 569
oxylipin 467
oxylipin signature 113

PAC libraries 421
palisade parenchyma formation 191
palmitoylation 429
petal 21
Petunia hybrida 725
Phaseolus lunatus 825
phloem 869
phloem proteins 735
PHO1 559
phosphate 383
phosphate starvation 559
phosphate translocators 285
phosphorylation 655
photomorphogenesis 345
photoperiod 591
photoreceptor 345
photosynthesis 383
phytochrome 883
pigment composition 191
PIP 231
plant cytoskeleton 859
plant cytosolic small Hsps 397
plant defense 837
plant development 613
plant disease resistance 569
plant DNA helicase 219
plant growth 613
plant mitochondria 637
plant–pathogen interactions 205
plasma membrane 79, 369, 775
plastid 241
plastid factor 523
plastid signal 883
pollen hydration 231
pollutant 467
poplar 11
Populus alba 147
potato 305
prenylated 775
prenylation 79
presequence 637
processing 45
programmed cell death 667
promoter 57
protein import 637
14-3-3 protein interaction 171
protein interaction 397
proteolysis 763
proton block 139

rapeseed 1
reactive oxygen species 667
receptor-like protein kinases 837
recombination 601
regulation 1, 171, 815
resistance 749
response regulator 703
reversible phosphorylation 171
riboregulators 559
rice 11
root 457
Rop GTPase 79

salicylic acid 837
salinity stress 511
SaMADS A 103
secologanin 797
secondary cell wall 477
secretory tissue 241
seed storage proteins 825
seedling growth 849
sex chromosomes 421
SH4 domain 429
shade avoidance 345
shoot apical meristem 91
signal transduction 317, 569, 693
Sinapis alba 103
sodium/myo-inositol symport 511
solanaceous plants 275
Solanum lycopersicum 345
splicing 317
starch 33
starch metabolism 285
stigma papillae 231
stomata 297
storage proteins 33
stress resistance 437
stress stimuli 785
stress tolerance 191
subcellular targeting 645
sucrose 147
sucrose synthesis 383
sucrose transport 503
sucrose transporter 869
sugar response 849
sugar sensing 869
susceptibility 749
syntaxin 369

targeting 429, 775
T-DNA 533
tetrapyrrole 883
threonine 357
tissue-specific expression 797
tobacco 583, 591, 763
tobacco protoplasts 397
tomato 317
tomato (*Lycopersicon esculentum*) 113
tonoplast 645
topoisomerase I 219
touch 655
transcript leader 45
transcription 57
transcription factor 275, 703
transgene integration 713
transgenic 305
transgenic plants 191, 265
transgenic wheat 713
translation 583
translation initiation factor 219
translational regulation 805
translocation intermediate 637
transposon 815
trichome 457
tryptophan 327
tunicamycin 825
two-component pseudo-response regulator 679
two-component regulatory system 703

ubiquitin 763
unwinding enzyme 219
uORF 45

vascular bundles 113
vegetative phase change 91
Vicia faba 297
viral movement protein 735
vitamin H 503
voltage independent 139

wheat 805, 895
wounding 655
WRKY proteins 837

Xanthomonas campestris pv. *campestris* 749
xylem 477
xyloglucan 147

Y-chromosome-specific clone 421
yeast complementation 489
yeast two-hybrid system 397, 679

zinc finger protein 191

THE plant JOURNAL

VOLUME 24

Editors

Thomas Altmann
Dianna Bowles
Clint Chapple
Jeff Dangl
Jean Dénarié
Jérôme Giraudat
Harry Klee
Maarten Koornneef
Cathie Martin
Gregory Martin
Julian Schroeder
Kazuo Shinozaki
Mark Stitt

Published by Blackwell Science Ltd,
in association with the Society for Experimental Biology

© 2000 Blackwell Science Ltd and the Society for Experimental Biology. Authorization to photocopy for internal or personal use or the internal or personal use of specific clients is granted by Blackwell Science Ltd and the Society for Experimental Biology for libraries and other users registered with the Copyright Clearance Centre (CCC) Transactional Reporting Service, provided that the base fee of $15.00 per copy is paid directly to the CCC, 222 Rosewood Drive, Suite 910, Danvers, MA 01923, USA. This consent does not extend to other kinds of copying, such as copying for general distribution for advertising or promotional purposes, for creating new collective works or for resale. Special requests should be addressed to the Chief Editor.
0960-7412/00 $15.00.

Published by Blackwell Science Ltd, in association with the Society for Experimental Biology

Blackwell Science Ltd

Osney Mead, Oxford OX2 0EL, UK
25 John Street, London WC1N 2BL, UK
23 Ainslie Place, Edinburgh EH3 6AJ, UK

Blackwell Science Inc., Commerce Place, 350 Main Street, Malden, MA 02148-5018, USA
Blackwell Science Asia, 54 University Street, Carlton South, Victoria 3053, Australia
Blackwell Wissenschafts-Verlag GmbH, Kurfürstendamm 57, 10707 Berlin, Germany
Blackwell Science KK, MG-Kodenmacho Building, 7–10 Kodenmacho Nihombashi, Cho-ku, Tokyo 103-0001, Japan
Blackwell Science Ltd, 10 rue Casimir Delavigne, 75006 Paris, France

ISSN 0960-7412

Typeset and printed by Information Press Ltd, Oxford, UK.

Contents

Number 1 October 2000

1 **Knockout of the regulatory site of 3-ketoacyl-ACP synthase III enhances short- and medium-chain acyl-ACP synthesis.** *A. Abbadi, M. Brummel and F. Spener*

11 **Activation of CDK-activating kinase is dependent on interaction with H-type cyclins in plants.** *M. Yamaguchi, T. Fabian, M. Sauter, R.P. Bhalerao, J. Schrader, G. Sandberg, M. Umeda and H. Uchimiya*

21 ***FRL1* is required for petal and sepal development in *Arabidopsis*.** *Y. Hase, A. Tanaka, T. Baba and H. Watanabe*

33 **Antisense-inhibition of ADP-glucose pyrophosphorylase in developing seeds of *Vicia narbonensis* moderately decreases starch but increases protein content and affects seed maturation.** *H. Weber, H. Rolletschek, U. Heim, S. Golombek, S. Gubatz and U. Wobus*

45 **Characterization and translational regulation of the arginine decarboxylase gene in carnation (*Dianthus caryophyllus* L.).** *K.S. Chang, S.H. Lee, S.B. Hwang and K.Y. Park*

57 **Transactivation of the *Brassica napus* napin promoter by ABI3 requires interaction of the conserved B2 and B3 domains of ABI3 with different *cis*-elements: B2 mediates activation through an ABRE, whereas B3 interacts with an RY/G-box.** *I. Ezcurra, P. Wycliffe, L. Nehlin, M. Ellerström and L. Rask*

67 ***Arabidopsis* DNA ligase IV is induced by γ-irradiation and interacts with an *Arabidopsis* homologue of the double strand break repair protein XRCC4.** *C.E. West, W.M. Waterworth, Q. Jiang and C.M. Bray*

79 **Maize ROP7 GTPase contains a unique, CaaX box-independent plasma membrane targeting signal.** *M. Ivanchenko, Z. Vejlupkova, R.S. Quatrano and J.E. Fowler*

91 **Mutations in the *WUSCHEL* gene of *Arabidopsis thaliana* result in the development of shoot without juvenile leaves.** *S. Hamada, H. Onouchi, H. Tanaka, M. Kudo, Y.-G. Liu, D. Shibata, C. Machida and Y. Machida*

103 **Cytokinin and gibberellin activate *SaMADS A*, a gene apparently involved in the regulation of the floral transition in *Sinapis alba*.** *F. Bonhomme, B. Kurz, S. Melzer, G. Bernier and A. Jacqmard*

113 **Tissue-specific oxylipin signature of tomato flowers: allene oxide cyclase is highly expressed in distinct flower organs and vascular bundles.** *B. Hause, I. Stenzel, O. Miersch, H. Maucher, R. Kramell, J. Ziegler and C. Wasternack*

127 TECHNICAL ADVANCE
Identification of plant actin-binding proteins by F-actin affinity chromatography. *S. Hu, S.R. Brady, D.R. Kovar, C.J. Staiger, G.B. Clark, S.J. Roux and G.K. Muday*

Number 2 October 2000

139 **Differential expression and regulation of K^+ channels in the maize coleoptile: molecular and biophysical analysis of cells isolated from cortex and vasculature.** *C.S. Bauer, S. Hoth, K. Haga, K. Philippar, N. Aoki and R. Hedrich*

147 **Evidence that endo-1,4-β-glucanases act on cellulose in suspension-cultured poplar cells.** *Y. Ohmiya, M. Samejima, M. Shiroishi, Y. Amano, T. Kanda, F. Sakai and T. Hayashi*

159 **The hormonal regulation of axillary bud growth in *Arabidopsis*.** *S.P. Chatfield, P. Stirnberg, B.G. Forde and O. Leyser*

171 **Post-translational regulation of cytosolic glutamine synthetase by reversible phosphorylation and 14-3-3 protein interaction.** *J. Finnemann and J.K. Schjoerring*

183 **Elevated levels of intrachromosomal homologous recombination in *Arabidopsis* overexpressing the *MIM* gene.** *M. Hanin, T. Mengiste, A. Bogucki and J. Paszkowski*

191 **A zinc finger protein RHL41 mediates the light acclimatization response in *Arabidopsis*.** *A. Iida, T. Kazuoka, S. Torikai, H. Kikuchi and K. Oeda*

205 **Three unique mutants of *Arabidopsis* identify *eds* loci required for limiting growth of a biotrophic fungal pathogen.** *J. Dewdney, T.L. Reuber, M.C. Wildermuth, A. Devoto, J. Cui, L.M. Stutius, E.P. Drummond and F.M. Ausubel*

219 **A DNA helicase from *Pisum sativum* is homologous to translation initiation factor and stimulates topoisomerase I activity.** *X.H. Pham, M.K. Reddy, N.Z. Ehtesham, B. Matta and N. Tuteja*

231 **Aquaporin *PIP* genes are not expressed in the stigma papillae in *Brassica oleracea*.** *M. Marin-Olivier, T. Chevalier, I. Fobis-Loisy, C. Dumas and T. Gaude*

241 **Molecular cloning of geranyl diphosphate synthase and compartmentation of monoterpene synthesis in plant cells.** *F. Bouvier, C. Suire, A. d'Harlingue, R.A. Backhuas and B. Camara*

253 TECHNICAL ADVANCE
An efficient method for dispersing *Ds* elements in the barley genome as a tool for determining gene function. *T. Koprek, D. McElroy, J. Louwerse, R. Williams-Carrier and P.G. Lemaux*

265 TECHNICAL ADVANCE
An estrogen receptor-based transactivator XVE mediates highly inducible gene expression in transgenic plants. *J. Zuo, Q.-W. Niu and N.-H. Chua*

275 TECHNICAL ADVANCE
A functional cloning strategy, based on a binary PVX-expression vector, to isolate HR-inducing cDNAs of plant pathogens. *F.L.W. Takken, R. Luderer, S.H.E.J. Gabriëls, N. Westerink, R. Lu, P.J.G.M. de Wit and M.H.A.J. Joosten*

Number 3 November 2000

285 **Plastidic metabolite transporters and their physiological functions in the inducible crassulacean acid metabolism plant *Mesembryanthemum crystallinum*.** *R.E. Häusler, B. Baur, J. Scharte, T. Teichmann, M. Eicks, K.L. Fischer, U.-I. Flügge, S. Schubert, A. Weber and K. Fischer*

297 **Dynamics of ionic activities in the apoplast of the sub-stomatal cavity of intact *Vicia faba* leaves during stomatal closure evoked by ABA and darkness.** *H.H. Felle, S. Hanstein, R. Steinmeyer and R. Hedrich*

305 **A potato α-glucosidase gene encodes a glycoprotein-processing α-glucosidase II-like activity. Demonstration of enzyme activity and effects of down-regulation in transgenic plants.** *M.A. Taylor, H.A. Ross, D. McRae, D. Stewart , I. Roberts, G. Duncan, F. Wright, S. Millam and H.V. Davies*

317 **Alternative splicing of a novel diacylglycerol kinase in tomato leads to a calmodulin-binding isoform.** *W.A. Snedden and E. Blumwald*

327 **Indole-3-glycerol phosphate, a branchpoint of indole-3-acetic acid biosynthesis from tryptophan biosynthetic pathway in *Arabidopsis thaliana*.** *J. Ouyang, X. Shao and J. Li*

335 **The characterization of differential calcium signalling in tobacco guard cells.** *N.T. Wood, A.C. Allan, A. Haley, M. Viry-Moussaïd and A.J. Trewavas*

345 **Physiological interactions of phytochromes A, B1 and B2 in the control of development in tomato.** *J.L. Weller, M.E.L. Schreuder, H. Smith, M. Koornneef and R.E. Kendrick*

357 **Identification of an *Arabidopsis thaliana* mutant accumulating threonine resulting from mutation in a new dihydrodipicolinate synthase gene.** *C. Sarrobert, M.-C. Thibaud, P. Contard-David, S. Gineste, N. Bechtold, C. Robaglia and L. Nussaume*

369 **Localization and control of expression of Nt-Syr1, a tobacco snare protein.** *B. Leyman, D. Geelen and M.R. Blatt*

383 **The role of inorganic phosphate in the development of freezing tolerance and the acclimatization of photosynthesis to low temperature is revealed by the *pho* mutants of *Arabidopsis thaliana*.** *V. Hurry, A. Strand, R. Furbank and M. Stitt*

397 **Transient expression and heat-stress-induced co-aggregation of endogenous and heterologous small heat-stress proteins in tobacco protoplasts.** *M. Kirschner, S. Winkelhaus, J.M. Thierfelder and L. Nover*

413 SHORT COMMUNICATION
Metabolic engineering of beta-carotene and lycopene content in tomato fruit. *C. Rosati, R. Aquilani, S. Dharmapuri, P. Pallara, C. Marusic, R. Tavazza, F. Bouvier, B. Camara and G. Giuliano*

421 TECHNICAL ADVANCE
Construction of male and female PAC genomic libraries suitable for identification of Y-chromosome-specific clones from the liverwort, *Marchantia polymorpha*. *S. Okada, M. Fujisawa, T. Sone, S. Nakayama, R. Nishiyama, M. Takenaka, S. Yamaoka, M. Sakaida, K. Kono, M. Takahama, K.T. Yamato, H. Fukuzawa, A. Brennicke and K. Ohyama*

Number 4 November 2000

429 **Membrane localization of a rice calcium-dependent protein kinase (CDPK) is mediated by myristoylation and palmitoylation.** *M.L. Martín and L. Busconi*

437 **A novel aldose/aldehyde reductase protects transgenic plants against lipid peroxidation under chemical and drought stresses.** *A. Oberschall, M. Deák, K. Török, L. Sass, I. Vass, I. Kovács, A. Fehér, D. Dudits and G.V. Horváth*

447 **Cloning and functional analysis of chicory root fructan 1-exohydrolase I (1-FEH I): a vacuolar enzyme derived from a cell-wall invertase ancestor? Mass fingerprint of the 1-FEH I enzyme.** *W. Van den Ende, A. Michiels, J. De Roover, P. Verhaert and A. Van Laere*

457 **MADS-box gene evolution beyond flowers: expression in pollen, endosperm, guard cells, roots and trichomes.** *E.R. Alvarez-Buylla, S.J. Liljegren, S. Pelaz, S.E. Gold, C. Burgeff, G.S. Ditta, F. Vergara-Silva and M.F. Yanofsky*

467 **Fatty acid ketodienes and fatty acid ketotrienes: Michael addition acceptors that accumulate in wounded and diseased *Arabidopsis* leaves.** *S. Vollenweider, H. Weber, S. Stolz, A. Chételat and E.E. Farmer*

477 **The *gapped xylem* mutant identifies a common regulatory step in secondary cell wall deposition.** *S.R. Turner and M. Hall*

489 **A member of a novel *Arabidopsis thaliana* gene family of candidate Mg^{2+} ion transporters complements a yeast mitochondrial group II intron-splicing mutant.** *I. Schock, J. Gregan, S. Steinhauser, R. Schweyen, A. Brennicke and V. Knoop*

503 **Plant sucrose-H^+ symporters mediate the transport of vitamin H.** *A. Ludwig, J. Stolz and N. Sauer*

511 **Na^+/*myo*-inositol symporters and Na^+/H^+-antiport in *Mesembryanthemum crystallinum*.** *S. Chauhan, N. Forsthoefel, Y. Ran, F. Quigley, D.E. Nelson and H.J. Bohnert*

523 **Chloroplast signalling in the light induction of nuclear *HSP70* genes requires the accumulation of chlorophyll precursors and their accessibility to cytoplasm/nucleus.** *J. Kropat, U. Oster, W. Rüdiger and C.F. Beck*

533 **Expression of a truncated tobacco *NtCBP4* channel in transgenic plants and disruption of the homologous *Arabidopsis CNGC1* gene confer Pb^{2+} tolerance.** *R. Sunkar, B. Kaplan, N. Bouché, T. Arazi, D. Dolev, I.N. Talke, F.J.M. Maathuis, D. Sanders, D. Bouchez and H. Fromm*

543 TECHNICAL ADVANCE
An aniline blue staining procedure for confocal microscopy and 3D imaging of normal and perturbed cellular phenotypes in mature *Arabidopsis* embryos. *S. Bougourd, J. Marrison and J. Haseloff*

551 TECHNICAL ADVANCE
Application of high-performance liquid chromatography with photodiode array detection to the metabolic profiling of plant isoprenoids. *P.D. Fraser, M.E.S. Pinto, D.E. Holloway and P.M. Bramley*

Number 5 December 2000

559 **Influence of cytokinins on the expression of phosphate starvation responsive genes in *Arabidopsis*.** *A.C. Martin, J.C. del Pozo, J. Iglesias, V. Rubio, R. Solano, A. de la Peña, A. Leyva and J. Paz-Ares*

569 **Oxidative burst and cognate redox signalling reported by luciferase imaging: identification of a signal network that functions independently of ethylene, SA and Me-JA but is dependent on MAPKK activity.** *J.J. Grant, B.-W. Yun and G.J. Loake*

583 **Functional characterization of the EMCV IRES in plants.** *P. Urwin, L. Yi, H. Martin, H. Atkinson and P.M. Gilmartin*

591 **A MADS domain gene involved in the transition to flowering in *Arabidopsis*.** *R. Borner, G. Kampmann, J. Chandler, R. Gleißner, E. Wisman, K. Apel and S. Melzer*

601 **A new hyperrecombinogenic mutant of *Nicotiana tabacum*.** *V. Gorbunova, N. Avivi-Ragolski, G. Shalev, I. Kovalchuk, S. Abbo, B. Hohn and A.A. Levy*

613 **Expression of the plant cyclin-dependent kinase inhibitor ICK1 affects cell division, plant growth and morphology.** *H. Wang, Y. Zhou, S. Gilmer, S. Whitwill and L.C. Fowke*

625 **AtFim1 is an actin filament crosslinking protein from *Arabidopsis thaliana*.** *D.R. Kovar, C.J. Staiger, E.A. Weaver and D.W. McCurdy*

637 **Studies on the topology of the protein import channel in relation to the plant mitochondrial processing peptidase integrated into the cytochrome bc_1 complex.** *P. Dessi, C. Rudhe and E. Glaser*

645 **Tonoplast subcellular localization of maize cytochrome b_5 reductases.** *P. Bagnaresi, D. Mazars-Marty, P. Pupillo, F. Marty and J.-F. Briat*

655 **Various abiotic stresses rapidly activate *Arabidopsis* MAP kinases ATMPK4 and ATMPK6.** *K. Ichimura, T. Mizoguchi, R. Yoshida, T. Yuasa and K. Shinozaki*

667 **NO way back: nitric oxide and programmed cell death in *Arabidopsis thaliana* suspension cultures.** *A. Clarke, R. Desikan, R.D. Hurst, J.T. Hancock and S.J. Neill*

679 **A stress-induced calcium-dependent protein kinase from *Mesembryanthemum crystallinum* phosphorylates a two-component pseudo-response regulator.** *O.R. Patharkar and J.C. Cushman*

693 **Promotive effect of brassinosteroids on cell division involves a distinct *CycD3*-induction pathway in *Arabidopsis*.** *Y. Hu, F. Bao and J. Li*

Number 6 December 2000

iii **Editorial**

703 ***Arabidopsis* ARR1 and ARR2 response regulators operate as transcriptional activators.** *H. Sakai, T. Aoyama and A. Oka*

713 **Widely separated multiple transgene integration sites in wheat chromosomes are brought together at interphase.** *R. Abranches, A.P. Santos, E. Wegel, S. Williams, A. Castilho, P. Christou, P. Shaw and E. Stoger*

725 ***NEC1*, a novel gene, highly expressed in nectary tissue of *Petunia hybrida*.** *Y.-X. Ge, G.C. Angenent, P.E. Wittich, J. Peters, J. Franken, M. Busscher, L.-M. Zhang, E. Dahlhaus, M.M. Kater, G.J. Wullems and T. Creemers-Molenaar*

735 **Proteolytic processing of CmPP36, a protein from the cytochrome b_5 reductase family, is required for entry into the phloem translocation pathway.** *B. Xoconostle-Cázares, R. Ruiz-Medrano and W.J. Lucas*

749 ***hxc2*, an *Arabidopsis* mutant with an altered hypersensitive response to *Xanthomonas campestris* pv. *campestris*.** *F. Godard, M.Lummerzheim, P. Saindrenan, C. Balagué and D. Roby*

763 **Cell cycle-dependent proteolysis and ectopic overexpression of cyclin B1 in tobacco BY2 cells.** *M.C. Criqui, Y. Parmentier, A. Derevier, W.-H. Shen, A. Dong and P. Genschik*

775 **Carboxyl-methylation of prenylated calmodulin CaM53 is required for efficient plasma membrane targeting of the protein.** *M. Rodríguez-Concepción, G. Toledo-Ortiz, S. Yalovsky, D. Caldelari and W. Gruissem*

785 **Auxin induces mitogenic activated protein kinase (MAPK) activation in roots of Arabidopsis seedlings.** *K. Mockaitis and S.H. Howell*

797 **Indole alkaloid biosynthesis in *Catharanthus roseus*: new enzyme activities and identification of cytochrome P450 CYP72A1 as secologanin synthase.** *S. Irmler, G. Schröder, B. St-Pierre, N.P. Crouch, M. Hotze, J. Schmidt, D. Strack, U. Matern and J. Schröder*

805 **Temperature-dependent translational regulation of the ER ω-3 fatty acid desaturase gene in wheat root tips.** *G. Horiguchi, T. Fuse, N. Kawakami, H. Kodama and K. Iba*

815 **The maize *LAG1-O* mutant suggests that reproductive cell lineages show unique gene expression patterns early in vegetative development.** *C.D. Giedt and C.F. Weil*

825 **Misfolding and aggregation of vacuolar glycoproteins in plant cells.** *F. Sparvoli, F. Faoro, M.G. Daminati, A. Ceriotti and R. Bollini*

837 **Identification of genes encoding receptor-like protein kinases as possible targets of pathogen- and salicylic acid-induced WRKY DNA-binding proteins in *Arabidopsis*.** *L. Du and Z. Chen*

849 **Monosaccharide/proton symporter AtSTP1 plays a major role in uptake and response of *Arabidopsis* seeds and seedlings to sugars.** *S.M. Sherson, G. Hemmann, G. Wallace, S. Forbes, V. Germain, R. Stadler, N. Bechtold, N. Sauer and S.M. Smith*

859 **Two kinesin-related proteins associated with the cold-stable cytoskeleton of carrot cells: characterization of a novel kinesin, DcKRP120-2.** *C. Barroso, J. Chan, V. Allan, J. Doonan, P. Hussey and C. Lloyd*

869 ***AtSUC3*, a gene encoding a new *Arabidopsis* sucrose transporter, is expressed in cells adjacent to the vascular tissue and in a carpel cell layer.** *S. Meyer, M. Melzer, E. Truernit, C. Hümmer, R. Besenbeck, R. Stadler and N. Sauer*

883 **Interactions between *hy1* and *gun* mutants of *Arabidopsis*, and their implications for plastid/nuclear signalling.** *G. Vinti, A. Hills, S. Campbell, J.R. Bowyer, N. Mochizuki, J. Chory and E. López-Juez*

895 TECHNICAL ADVANCE
Double-stranded RNA interferes with gene function at the single-cell level in cereals. *P. Schweizer, J. Pokorny, P. Schulze-Lefert and R. Dudler*

905 **Author Index**

911 **Subject Index**